

**EECS 210**  
 Fall 2006  
 Tu, Th 12:30-2  
 400 Cory

**Applied Electromagnetic Theory**  
 Office Hours  
 M, (W), 11AM  
 Tu, Th, (F) 10AM

Prof. A. R. Neureuther,  
 509 Cory Hall, 2-4590  
 neureuth@eecs



### Homework # 3: Due start of class Th Sep 17th

3.1) **Integral Equations by Hand and Cylindrical Geometry Practice:** Consider two small grounded conducting circular cylinders of radius  $a = 0.1\text{m}$  with centers at  $(x,y) = (0,0)$  for cylinder 1 and  $(x,y) = (1,0)$  for cylinder 2. A line charge of strength  $\lambda_{S1}$  C/m is brought into position  $(0, 0.5)$ . Since  $a$  is much smaller than the separation between the charges and cylinders the charge induced on the cylinder can be modeled as a line charge  $\lambda_1$  and  $\lambda_2$  at the center of the cylinder

- Write down the integral representation for the potential at a location  $(x,y)$  for  $\lambda_{S1}$  in the presence of a conducting cylinder at voltage  $V$  of arbitrary shape the Green's function for a line charge in free space.
- Show how this integral representation specializes to multiple circular cylinder conductors that are grounded and for which the induced charge can be located at the center of the cylinder. Hint: For the self contribution to the potential use the uniform electric field on a circular cylinder of radius  $a$ .
- Generate a two-by-two matrix equation  $\bar{\Phi} = \bar{0} = \bar{A}q + \bar{B}$  for the zero potential at observation points on the two circular cylinders as a function of the potential contributed by the induced charges on the two circular cylinders (that is the  $Aq$  term) and the contribution to the potential on each of the cylinders from the source  $q_{S1}$  (that is the  $B$  term).
- Solve the matrix equation for the induced charges.
- Compute the potential at location  $(2,0)$ .
- Repeat for a source of strength  $q_{S2}$  at location  $(2,0)$  and find the potential at  $(0,0.5)$ . Show that reciprocity holds  $q_{S1}\Phi(2,0) = q_{S2}\Phi(0,0.5)$  even though the induced charges on the cylinder appear to be unrelated.

3.2) **Creating Ortho-Normal Functions:** A power series expansion  $y(x) = 1 + 0.5x$  on the interval  $[0,1]$  is to be converted to an expansion in orthonormal functions

$y(x) = B_0\phi_0(x) + B_1\phi_1(x)$  where the first function is constant  $\phi_0 = C_{00}$  and the second is linear  $\phi_1 = C_{10} + C_{11}x$ .

- Find  $C$  by applying normalization to  $\phi_0$ .
- Use orthogonality to find one condition on  $C_{10}$  and  $C_{11}$ .
- Use normalization to find a second constraint on  $C_{10}$  and  $C_{11}$  and solve for  $\phi_1$ .

d) Compute the mean squared value of  $y(x)$  directly and check that it equals  $MS = \sum_{n=0}^1 B_n^2$ .

e) Suppose  $n$  were to go to 4 in defining orthonormal functions. Determine the number of unknown coefficients  $C$  that would have to be determined. Then show that the number of constraint equations from orthogonality plus the number from normalization give this number.

3.3) **Constancy of the product of spatial width and Fourier Integral width:** For the definition of the Fourier transform in Jackson on pp. 69 take the product of the full-width half-maximum of  $f(x)$  with the full-width half-maximum  $A(k)$  for a) rectangular shape, b) triangular shape, and c) Gaussian. Are they approximately the same?