

CS-184: Computer Graphics


Lecture #15: Natural Splines, B-Splines, and NURBS

Prof. James O'Brien
University of California, Berkeley

V2006S-15-0.9

Natural Splines

- Draw a “smooth” line through several points



A real draftsman's spline.


Image from Carl de Boor's webpage.

Natural Cubic Splines

- Given $n + 1$ points
 - Generate a curve with n segments
 - Curves passes through points
 - Curve is C^2 continuous
- Use cubics because lower order is better...

3


Natural Cubic Splines



$$\mathbf{x}(u) = \begin{cases} \mathbf{s}_1(u) & \text{if } 0 \leq u < 1 \\ \mathbf{s}_2(u - 1) & \text{if } 1 \leq u < 2 \\ \mathbf{s}_3(u - 2) & \text{if } 2 \leq u < 3 \\ \vdots & \\ \mathbf{s}_n(u - (n - 1)) & \text{if } n - 1 \leq u \leq n \end{cases}$$

4

Natural Cubic Splines



$$\begin{array}{lll} s_i(0) = p_{i-1} & i = 1 \dots n & \leftarrow n \text{ constraints} \\ s_i(1) = p_i & i = 1 \dots n & \leftarrow n \text{ constraints} \end{array}$$

$$\begin{array}{lll} s'_i(1) = s'_{i+1}(0) & i = 1 \dots n-1 & \leftarrow n-1 \text{ constraints} \\ s''_i(1) = s''_{i+1}(0) & i = 1 \dots n-1 & \leftarrow n-1 \text{ constraints} \end{array}$$

$$s''_1(0) = s''_n(1) = 0 \qquad \qquad \qquad \leftarrow 2 \text{ constraints}$$

Total $4n$ constraints

5

Natural Cubic Splines

- Interpolate data points
- No convex hull property
- Non-local support
 - Consider matrix structure...
- C^2 using cubic polynomials


6

B-Splines

- Goal: C^2 cubic curves with local support
 - Give up interpolation
 - Get convex hull property
- Build basis by designing “hump” functions

7

B-Splines



$$b(u) = \begin{cases} b_{-2}(u) & \text{if } u_{-2} \leq u < u_{-1} \\ b_{-1}(u) & \text{if } u_{-1} \leq u < u_0 \\ b_0(u) & \text{if } u_0 \leq u < u_1 \\ b_1(u) & \text{if } u_1 \leq u < u_2 \\ b_2(u) & \text{if } u_2 \leq u \leq u_{+2} \end{cases}$$


$$\begin{aligned} b''_{-2}(u_{-2}) &= b'_{-2}(u_{-2}) = b_{-2}(u_{-2}) = 0 & \leftarrow 3 \text{ constraints} \\ b''_{+2}(u_{+2}) &= b'_{+2}(u_{+2}) = b_{+2}(u_{+2}) = 0 & \leftarrow 3 \text{ constraints} \end{aligned}$$

$$\begin{aligned} b_{-2}(u_{-1}) &= b_{-1}(u_{-1}) \\ b_{-1}(u_0) &= b_{+1}(u_0) \\ b_{+1}(u_{+1}) &= b_{+2}(u_{+1}) \end{aligned} \quad \leftarrow \begin{array}{l} \text{Repeat for } b' \text{ and } b'' \\ 3 \times 3 = 9 \text{ constraints} \end{array}$$

Total 15 constraints need one more

8

B-Splines



$$b(u) = \begin{cases} b_{-2}(u) & \text{if } u_{-2} \leq u < u_{-1} \\ b_{-1}(u) & \text{if } u_{-1} \leq u < u_0 \\ b_0(u) & \text{if } u_0 \leq u < u_1 \\ b_1(u) & \text{if } u_1 \leq u < u_2 \\ b_2(u) & \text{if } u_2 \leq u \leq u_+ \end{cases}$$

$$\begin{aligned} b''_{-2}(u_{-2}) &= b'_{-2}(u_{-2}) = b_{-2}(u_{-2}) = 0 & \leftarrow 3 \text{ constraints} \\ b''_{+2}(u_2) &= b'_{+2}(u_2) = b_{+2}(u_2) = 0 & \leftarrow 3 \text{ constraints} \end{aligned}$$

$$\begin{aligned} b_{-2}(u_{-1}) &= b_{-1}(u_{-1}) \\ b_{-1}(u_0) &= b_0(u_0) \\ b_0(u_1) &= b_1(u_1) \end{aligned} \quad \leftarrow \begin{array}{l} \text{Repeat for } b' \text{ and } b'' \\ 3 \times 3 = 9 \text{ constraints} \end{array}$$

$$b_{-2}(u_{-2}) + b_{-1}(u_{-1}) + b_0(u_0) + b_1(u_1) = 1 \leftarrow 1 \text{ constraint (convex hull)}$$

Total 16 constraints

9

B-Splines

- Build a curve w/ overlapping bumps
- Continuity
 - Inside bumps C^2
 - Bumps “fade out” with C^2 continuity
- Boundaries
 - Circular
 - Repeat end points
 - Extra end points

10

B-Splines

- Notation

- The basis functions are the $b_i(u)$
- “Hump” functions are the concatenated function
 - Sometimes the humps are called basis... can be confusing
- The u_i are the knot locations
- The weights on the hump/basis functions are control points

11

B-Splines

- Similar construction method can give higher continuity with higher degree polynomials
- Repeating knots drops continuity
 - Limit as knots approach each other
- Still cubics, so conversion to other cubic basis is just a matrix multiplication


12

B-Splines

- Geometric construction
 - Due to Cox and de Boor
 - My own notation, beware if you compare w/ text
- Let hump centered on u_i be $N_{i,4}(u)$
Cubic is order 4
 $N_{i,k}(u)$ is order k hump, centered at u_i
Note: i is integer if k is even
else $(i + 1/2)$ is integer

13

B-Splines

$$N_{i,1}(u) = \begin{cases} 1 & \text{if } u_{i-\frac{1}{2}} \leq u < u_{i+\frac{1}{2}} \\ 0 & \text{else} \end{cases}$$

$$N_{i,k}(u) = \frac{(u - u_{i-k/2}) N_{i-k/2, k-1}(u)}{u_{i+k/2-1} - u_{i-k/2}}$$


↑ Term #1

$$\frac{(u_{i+k/2} - u) N_{i+k/2, k-1}(u)}{u_{i+k/2} - u_{i-k/2+1}}$$

Term #2

Recursive defn.

14



15

NURBS

- **Nonuniform Rational B-Splines**
 - Basically B-Splines using homogeneous coordinates
 - Transform under perspective projection
 - A bit of extra control

16

NUBRS

$$\mathbf{p}_i = \begin{bmatrix} p_{ix} \\ p_{iy} \\ p_{iz} \\ p_{iw} \end{bmatrix} \quad \mathbf{x}(u) = \frac{\sum_i \begin{bmatrix} p_{ix} \\ p_{iy} \\ p_{iz} \end{bmatrix} N_i(u)}{\sum_i p_{iw} N_i(u)}$$

- Non-linear in the control points
- The p_{iw} are sometimes called “weights”