
1 Markov Chain Terminology

In this question, we will walk you through terms related to Markov chains. Keep in mind the following theorems and ideas as you work.

- 1. (Irreducibility) A Markov chain is irreducible if, starting from any state i, the chain can transition to any other state j, possibly in multiple steps.
- 2. (Periodicity) $d(i) := \gcd\{n > 0 \mid P^n(i,i) = \Pr[X_n = i \mid X_0 = i] > 0\}, i \in \mathcal{X}$. If $d(i) = 1 \ \forall i \in \mathcal{X}$, then the Markov chain is aperiodic; otherwise it is periodic.
- 3. (Matrix Representation) Define the transition probability matrix P by filling entry (i, j) with probability P(i, j).
- 4. (Invariance) A distribution π is invariant for the transition probability matrix P if it satisfies the following balance equations: $\pi = \pi P$.

Use the above theorems and the Markov chain below to answer the following questions.



- (a) For what values of a and b is the above Markov chain irreducible? Reducible?
- (b) For a = 1, b = 1, prove that the above Markov chain is periodic.
- (c) For 0 < a < 1, 0 < b < 1, prove that the above Markov chain is aperiodic.
- (d) Construct a transition probability matrix using the above Markov chain.
- (e) Write down the balance equations for this Markov chain and solve them. Assume that the Markov chain is irreducible.

2 Allen's Umbrellas

Every morning, Allen walks from his home to Soda, and every evening, Allen walks from Soda to his home. Suppose that Allen has two umbrellas in his possession, but he sometimes leaves his umbrellas behind. Specifically, before leaving from his home or Soda, he checks the weather. If it is raining outside, he will bring his umbrella (that is, if there is an umbrella where he currently is). If it is not raining outside, he will forget to bring his umbrella. Assume that the probability of rain is *p*.

We will model this as a Markov chain. Let $\mathscr{X} = \{0,1,2\}$ be the set of states, where the state i represents the number of umbrellas in his current location. Write down the transition matrix, determine if the distribution of X_n converges to the invariant distribution, and compute the invariant distribution. Determine the long-term fraction of time that Allen will walk through rain with no umbrella.

3 Continuous Intro

(a) Is

$$f(x) = \begin{cases} 2x, & 0 \le x \le 1\\ 0, & \text{otherwise} \end{cases}$$

a valid density function? Why or why not? Is it a valid CDF? Why or why not?

(b) Calculate $\mathbf{E}[X]$ and var(X) for X with the density function

$$f(x) = \begin{cases} 1/\ell, & 0 \le x \le \ell, \\ 0, & \text{otherwise.} \end{cases}$$

(c) Suppose *X* and *Y* are independent and have densities

$$f_X(x) = \begin{cases} 2x, & 0 \le x \le 1, \\ 0, & \text{otherwise,} \end{cases}$$
$$f_Y(y) = \begin{cases} 1, & 0 \le y \le 1, \\ 0, & \text{otherwise.} \end{cases}$$

What is their joint distribution?

(d) Calculate $\mathbf{E}[XY]$ for the above X and Y.

4 Uniform Distribution

You have two spinners, each having a circumference of 10, with values in the range [0, 10). If you spin both (independently) and let X be the position of the first spinner and Y be the position of the second spinner, what is the probability that X > 5, given that Y > X?